

On Complex (non analytic) Chebyshev Polynomials in $\mathbb{C}^2$

I. Moale* and P. Yuditskii†

February 10, 2010

Dedicated to the memory of Franz Peherstorfer

Abstract

We consider the problem of finding a best uniform approximation to the standard monomial on the unit ball in $\mathbb{C}^2$ by polynomials of lower degree with complex coefficients. We reduce the problem to a one-dimensional weighted minimization problem on an interval. In a sense, the corresponding extremal polynomials are uniform counterparts of the classical orthogonal Jacobi polynomials. They can be represented by means of special conformal mappings on the so-called comb-like domains. In these terms, the value of the minimal deviation and the representation for a polynomial of best approximation for the original problem are given. Furthermore, we derive asymptotics for the minimal deviation.

1 Introduction

We consider the standard basis in the set of (non analytic) complex polynomials in $\mathbb{C}^2$:

$$\{z_1^{k_1} \bar{z}_1^{l_1} z_2^{k_2} \bar{z}_2^{l_2}\}_{k_1 \geq 0, l_1 \geq 0, k_2 \geq 0, l_2 \geq 0}, \quad (z_1, z_2) \in \mathbb{C}^2.$$

As usual $k_1 + l_1 + k_2 + l_2$ is called the total degree of the given monomial.

In what follows we use the following notations: Π_n denotes the set of polynomials with complex coefficients of total degree less or equal n , and $\|P\|$ denotes the uniform norm of $P \in \Pi_n$ in the complex ball

$$\|P\| = \sup_{(z_1, z_2) \in \mathbb{B}} |P(z_1, z_2)|, \quad \mathbb{B} = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^2 \leq 1\}.$$

*Supported by the Austrian Science Fund FWF, project number: P20413-N18

†Supported by the Austrian Science Fund FWF, project number: P22025-N18




Figure 1: Domain $\Omega_2(\alpha, \beta)$

Analogously to the classical Chebyshev polynomial, we consider the best approximation on the ball $\mathbb{B}$ of the monomial $z_1^{k_1} \bar{z}_1^{l_1} z_2^{k_2} \bar{z}_2^{l_2}$ by polynomials of total degree less than $n := k_1 + l_1 + k_2 + l_2$. Such a polynomial $\tilde{T}_{k_1, l_1, k_2, l_2}(z_1, z_2) = z_1^{k_1} \bar{z}_1^{l_1} z_2^{k_2} \bar{z}_2^{l_2} + \dots$, which we call a polynomial of least deviation from zero on $\mathbb{B}$, is not unique but the minimal deviation L_{k_1, l_1, k_2, l_2} is well defined,

$$L_{k_1, l_1, k_2, l_2} := \inf_{P \in \Pi_{n-1}} \|z_1^{k_1} \bar{z}_1^{l_1} z_2^{k_2} \bar{z}_2^{l_2} - P(z_1, z_2)\|.$$

It is convenient to work with the normalized polynomial $T_{k_1, l_1, k_2, l_2} := \tilde{T}_{k_1, l_1, k_2, l_2} / \|\tilde{T}_{k_1, l_1, k_2, l_2}\|$. Thus

$$T_{k_1, l_1, k_2, l_2}(z_1, z_2) = \Lambda_{k_1, l_1, k_2, l_2} z_1^{k_1} \bar{z}_1^{l_1} z_2^{k_2} \bar{z}_2^{l_2} + \dots,$$

where $\Lambda_{k_1, l_1, k_2, l_2} = 1/L_{k_1, l_1, k_2, l_2}$.

Concerning polynomials of least deviation from zero on the unit ball in $\mathbb{R}^2$, several approaches are known so far, see [4, 8, 13] and also [11]. Of foremost importance to us was the representation of the extremal polynomial given by Braß in [4]. Here we essentially simplify and generalize his construction.

In approximation theory, a special role of the conformal mappings on so-called comb-like domains is well known, see e.g. the book [1], the survey [14] and the references on original papers therein, in particular [2, 9]. For recent developments in this direction, see [6, 12]. By analogy with the MacLane-Vinberg special representation for polynomials and entire functions [10, 15], in this paper to nonnegative real numbers α, β and an integer $n \geq 0$ we associate a horizontal strip with $n + 1$ horizontal cuts, see Fig. 1:

$$\Omega_n(\alpha, \beta) = \{w = u + iv : -\beta < \frac{v}{\pi} < \alpha + n\} \setminus \bigcup_{j=0}^n \{w = u + iv : \frac{v}{\pi} = j, u \leq 0\}.$$

Note that the boundary of the domain contains $n + 3$ infinite points:

$$\begin{aligned}\infty_0 &= -\infty + iv, n < \frac{v}{\pi} < n + \alpha, & \infty_j &= -\infty + iv, n - j < \frac{v}{\pi} < n - j + 1, \\ \infty_{n+1} &= -\infty + iv, -\beta < \frac{v}{\pi} < 0, & \infty_{n+2} &= +\infty + iv, -\beta < \frac{v}{\pi} < \alpha + n.\end{aligned}$$

Let $w : \mathbb{C}_+ \rightarrow \Omega_n(\alpha, \beta)$ be the conformal mapping of the upper half-plane onto $\Omega_n(\alpha, \beta)$ with the following normalization

$$w(0) = \infty_0, \quad w(1) = \infty_{n+1}, \quad w(\infty) = \infty_{n+2}.$$

It is easy to see that it has the following asymptotics at infinity ($z \rightarrow \infty$)

$$w(z) = w_n(z; \alpha, \beta) = (\alpha + \beta + n) \ln z + C_n(\alpha, \beta) - \beta\pi i + O(1/z) \quad (1.1)$$

The real constant $C_n(\alpha, \beta)$ is uniquely defined by the domain (a kind of capacity).

Due to the evident symmetry

$$\Lambda_{k_1, l_1, k_2, l_2} = \Lambda_{l_1, k_1, k_2, l_2} = \Lambda_{k_1, l_1, l_2, k_2}$$

we can assume that $k_1 \geq l_1$ and $k_2 \geq l_2$. Our first result is

Theorem 1.1. *In the above introduced notations*

$$\ln \Lambda_{k_1, l_1, k_2, l_2} = C_{l_1 + l_2} \left(\frac{k_1 - l_1}{2}, \frac{k_2 - l_2}{2} \right).$$

Below we give the representation for a polynomial $T_{k_1, l_1, k_2, l_2}(z_1, z_2)$ of least deviation from zero. For this, we establish a connection between the conformal mapping $w_n(z; \alpha, \beta)$ and a weighted 1-D extremal problem on $[0, 1]$. In a sense, in the following proposition we define *uniform Jacobi polynomials*, compare to the classical orthogonal ones [3].

Proposition 1.2. *Let $\xi_j = w_n^{-1}(\infty_j; \alpha, \beta)$, $1 \leq j \leq n$. Then*

$$e^{w_n(t; \alpha, \beta)} = t^\alpha (1 - t)^\beta e^{C_n(\alpha, \beta)} (t - \xi_1) \dots (t - \xi_n), \quad (1.2)$$

where $\tilde{\mathcal{J}}_n(t; \alpha, \beta) := (t - \xi_1) \dots (t - \xi_n) = t^n + \dots$ is the polynomial of least deviation from zero on $[0, 1]$ with respect to the weight $t^\alpha (1 - t)^\beta$. Moreover

$$\|\tilde{\mathcal{J}}_n(t; \alpha, \beta)\| = \sup_{0 \leq t \leq 1} t^\alpha (1 - t)^\beta |\tilde{\mathcal{J}}_n(t; \alpha, \beta)| = e^{-C_n(\alpha, \beta)}, \quad (1.3)$$

that is,

$$e^{w_n(t; \alpha, \beta)} = t^\alpha (1 - t)^\beta \mathcal{J}_n(t; \alpha, \beta),$$

as before $\mathcal{J}_n(t; \alpha, \beta) := \tilde{\mathcal{J}}_n(t; \alpha, \beta) / \|\tilde{\mathcal{J}}_n(t; \alpha, \beta)\|$.

We point out that $\xi_l < \xi_{l+1}$ for all $1 \leq l \leq n-1$.

Theorem 1.3. *Let $\alpha = \frac{k_1-l_1}{2} \geq 0$ and $\beta = \frac{k_2-l_2}{2} \geq 0$. Let us factorize $\tilde{\mathcal{J}}_{l_1+l_2}(t; \alpha, \beta) = \tilde{\mathcal{J}}_{l_1}^{(1)}(t)\tilde{\mathcal{J}}_{l_2}^{(2)}(t)$ in polynomials of degrees l_1 and l_2 respectively in the following way*

$$\begin{aligned}\tilde{\mathcal{J}}_{l_1}^{(1)}(t) &= (t - \xi_1) \dots (t - \xi_{l_1}) \\ \tilde{\mathcal{J}}_{l_2}^{(2)}(t) &= (t - \xi_{l_1+1}) \dots (t - \xi_{l_1+l_2}).\end{aligned}\tag{1.4}$$

Then

$$T_{k_1, l_1, k_2, l_2}(z_1, z_2) = e^{C_{l_1+l_2}(\alpha, \beta)} z_1^{k_1-l_1} z_2^{k_2-l_2} \tilde{\mathcal{J}}_{l_1}^{(1)}(|z_1|^2) (-1)^{l_2} \tilde{\mathcal{J}}_{l_2}^{(2)}(1 - |z_2|^2)\tag{1.5}$$

Finally we present the following asymptotic relation for the value of the minimal deviation.

Theorem 1.4. *Assume that the following limits exist*

$$\varkappa_1 = \lim_{n \rightarrow \infty} \frac{k_1}{n}, \lambda_1 = \lim_{n \rightarrow \infty} \frac{l_1}{n}, \varkappa_2 = \lim_{n \rightarrow \infty} \frac{k_2}{n}, \lambda_2 = \lim_{n \rightarrow \infty} \frac{l_2}{n},$$

where $n = k_1 + l_1 + k_2 + l_2$. Then

$$\lim_{n \rightarrow \infty} L_{k_1, l_1, k_2, l_2}^{\frac{2}{n}} = (\lambda_1 + \lambda_2)^{\lambda_1 + \lambda_2} (\varkappa_1 + \varkappa_2)^{\varkappa_1 + \varkappa_2} (\lambda_1 + \varkappa_2)^{\lambda_1 + \varkappa_2} (\varkappa_1 + \lambda_2)^{\varkappa_1 + \lambda_2}.$$

2 Reduction to 1-D problem

First, we reduce our complex two-dimensional approximation problem to a weighted approximation problem in two real variables on the standard triangle $\Delta := \{(t_1, t_2) \in \mathbb{R}^2 : t_1 \geq 0, t_2 \geq 0, t_1 + t_2 \leq 1\}$.

For a continuous function f on Δ , we define $\|f\|_{\Delta} := \max_{(t_1, t_2) \in \Delta} |f(t_1, t_2)|$.

By $Y_{l_1, l_2}(t_1, t_2; \alpha, \beta) = M_{l_1, l_2}(\alpha, \beta) t_1^{l_1} t_2^{l_2} + \dots$ we denote a normalized polynomial of least deviation from zero on Δ with respect to the weight function $t_1^{\alpha} t_2^{\beta}$.

Proposition 2.1. *Let $\alpha = \frac{k_1-l_1}{2}$ and $\beta = \frac{k_2-l_2}{2}$. Then*

$$z_1^{k_1-l_1} z_2^{k_2-l_2} Y_{l_1, l_2}(|z_1|^2, |z_2|^2; \alpha, \beta)$$

is a normalized polynomial of least deviation from zero on $\mathbb{B}$, that is,

$$\Lambda_{k_1, l_1, k_2, l_2} = M_{l_1, l_2}(\alpha, \beta).$$

Proof. Let us remark that due to the symmetries of $\mathbb{B}$, if $T_{k_1, l_1, k_2, l_2}(z_1, z_2)$ is a polynomial of least deviation from zero, then for any $\theta_1, \theta_2 \in [0, 2\pi]$, the polynomials

$$T_{k_1, l_1, k_2, l_2}(e^{i\theta_1} z_1, e^{i\theta_2} z_2) e^{-i(k_1 - l_1)\theta_1} e^{-i(k_2 - l_2)\theta_2}$$

and

$$\frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} T_{k_1, l_1, k_2, l_2}(e^{i\theta_1} z_1, e^{i\theta_2} z_2) e^{-i(k_1 - l_1)\theta_1} e^{-i(k_2 - l_2)\theta_2} d\theta_1 d\theta_2 \quad (2.1)$$

are also polynomials of least deviation from zero.

It is easy to see that the polynomial in (2.1) is of the form

$$\begin{aligned} \Lambda_{k_1, l_1, k_2, l_2} z_1^{k_1 - l_1} z_2^{k_2 - l_2} \left[|z_1|^{2l_1} |z_2|^{2l_2} + \sum_{c_{j_1 + j_2} \leq l_1 + l_2 - 1} a_{j_1, j_2} |z_1|^{2j_1} |z_2|^{2j_2} \right] \\ =: \Lambda_{k_1, l_1, k_2, l_2} z_1^{k_1 - l_1} z_2^{k_2 - l_2} \tilde{P}_{l_1, l_2}(|z_1|^2, |z_2|^2) \end{aligned}$$

where $a_{j_1, j_2} \in \mathbb{C}$. Note that

$$\hat{T}_{k_1, l_1, k_2, l_2}(z_1, z_2) := z_1^{k_1 - l_1} z_2^{k_2 - l_2} Q_{l_1, l_2}(|z_1|^2, |z_2|^2),$$

where $Q_{l_1, l_2}(|z_1|^2, |z_2|^2) := \Lambda_{k_1, l_1, k_2, l_2} \text{Re} \tilde{P}_{l_1, l_2}(|z_1|^2, |z_2|^2)$, is still a normalized polynomial of least deviation from zero on $\mathbb{B}$.

Since $(z_1, z_2) \in \mathbb{B}$ is equivalent to $(t_1, t_2) \in \Delta$, where $t_1 := |z_1|^2$, $t_2 := |z_2|^2$, we have that

$$\|\hat{T}_{k_1, l_1, k_2, l_2}\|_{\mathbb{B}} = \|t_1^{(k_1 - l_1)/2} t_2^{(k_2 - l_2)/2} Q_{l_1, l_2}\|_{\Delta},$$

which gives the assertion. $\square$

Now we give a sufficient condition for a polynomial to be a weighted polynomial of least deviation from zero on Δ . In the next section we show the existence of a polynomial satisfying this condition.

As before $\mathcal{J}_n(t; \alpha, \beta)$ denotes the normalized polynomial of least deviation from zero on $[0, 1]$ with respect to the weight function $t^\alpha(1 - t)^\beta$.

Proposition 2.2. *Let $\alpha, \beta \geq 0$ and let $c_n = c_n(\alpha, \beta) > 0$ be the leading coefficient of $\mathcal{J}_n(t; \alpha, \beta)$, $\mathcal{J}_n(t; \alpha, \beta) = c_n t^n + \dots$. If there exists a polynomial*

$$P_{l_1, l_2}(t_1, t_2) = c_{l_1 + l_2}(\alpha, \beta) t_1^{l_1} t_2^{l_2} + \dots \text{ with } \|t_1^\alpha t_2^\beta P_{l_1, l_2}\|_{\Delta} \leq 1 \quad (2.2)$$

such that $P_{l_1, l_2}(t, 1-t) = (-1)^{l_2} \mathcal{J}_{l_1+l_2}(t; \alpha, \beta)$ for all $t \in [0, 1]$, then

$$M_{l_1, l_2}(\alpha, \beta) = c_{l_1+l_2}(\alpha, \beta). \quad (2.3)$$

That is, the given P_{l_1, l_2} is a normalized polynomial of least deviation from zero on Δ with respect to the weight $t_1^\alpha t_2^\beta$.

Proof. Actually, we have to prove (2.3).

By the fact that Y_{l_1, l_2} is a polynomial of least deviation from zero, from (2.2) we have immediately that $M_{l_1, l_2} \geq c_{l_1+l_2}$.

On the other hand, let us restrict Y_{l_1, l_2} to the line $t_1 = t, t_2 = 1-t$:

$$Q(t) := (-1)^{l_2} Y_{l_1, l_2}(t, 1-t) = M_{l_1, l_2} t^{l_1+l_2} + \dots$$

Since $|Q(t)t^\alpha(1-t)^\beta| \leq 1$ for all $t \in [0, 1]$, the extremal property of $\mathcal{J}_{l_1+l_2}$ implies $M_{l_1, l_2} \leq c_{l_1+l_2}$. Thus the statement is proved. $\square$

3 Proofs of Proposition 1.2, Theorems 1.1 and 1.3

Proof of Proposition 1.2. Let $\eta_k = w_n^{-1}((n-k)\pi i; \alpha, \beta)$, $0 \leq k \leq n$. From the Schwarz-Christoffel formula, see e.g. [5], we obtain the following expression for the differential of the conformal mapping $w_n(z; \alpha, \beta)$:

$$dw_n(z; \alpha, \beta) = C \frac{\prod_{k=0}^n (z - \eta_k)}{z(z-1) \prod_{j=1}^n (z - \xi_j)} dz, \quad (3.1)$$

where $C \in \mathbb{C}$ is a constant. Having in mind the asymptotic behavior at the infinite boundary points of the domain $\Omega_n(\alpha, \beta)$ we get the following expansion into partial fraction for (3.1):

$$dw_n(z; \alpha, \beta) = \left(\frac{\alpha}{z} + \frac{\beta}{z-1} + \sum_{j=1}^n \frac{1}{z - \xi_j} \right) dz.$$

Hence

$$w_n(z; \alpha, \beta) = \alpha \ln z + \beta \ln(z-1) + \sum_{j=1}^n \ln(z - \xi_j) + C_1, \quad (3.2)$$

where $C_1 \in \mathbb{C}$ is a constant. Relation (1.2) follows now immediately from (3.2).




Figure 2: Graph of $t^\alpha(1-t)^\beta \mathcal{J}_2(t; \alpha, \beta)$, $\alpha = 1/2, \beta = 2$.

From the boundary correspondence for the given conformal mapping we get that the function $t^\alpha(1-t)^\beta \mathcal{J}_n(t; \alpha, \beta)$ alternates $n+1$ times between ± 1 on $[0, 1]$, see Fig. 2. Thus the Chebyshev alternation theorem implies that $\mathcal{J}_n(t; \alpha, \beta)$ is indeed the polynomial of least deviation from zero with respect to the given weight with leading coefficient $c_n(\alpha, \beta) = e^{C_n(\alpha, \beta)}$. $\square$

Proof of Theorems 1.1 and 1.3. By Proposition 2.1, the assertion of Theorem 1.3 follows if we are able to show that

$$\begin{aligned} P_{l_1, l_2}(t_1, t_2; \alpha, \beta) &:= e^{C_{l_1+l_2}(\alpha, \beta)} \tilde{J}_{l_1}^{(1)}(t_1) (-1)^{l_2} \tilde{J}_{l_2}^{(2)}(1-t_2) \\ &= e^{C_{l_1+l_2}(\alpha, \beta)} t_1^{l_1} t_2^{l_2} + \dots \end{aligned} \quad (3.3)$$

is a normalized polynomial of least deviation from zero on Δ with respect to the weight $t_1^\alpha t_2^\beta$, for which we will use Proposition 2.2.

By restricting the polynomial P_{l_1, l_2} to the line $t_1 := t, t_2 := 1-t$, we obtain that for all $t \in [0, 1]$:

$$P_{l_1, l_2}(t, 1-t; \alpha, \beta) = (-1)^{l_2} \mathcal{J}_{l_1+l_2}(t; \alpha, \beta). \quad (3.4)$$

Thus it remains to show that $\|t_1^\alpha t_2^\beta P_{l_1, l_2}\|_\Delta \leq 1$.

Let

$$F(t_1, t_2; \alpha, \beta) := t_1^\alpha t_2^\beta P_{l_1, l_2}(t_1, t_2; \alpha, \beta).$$

Clearly $F(t_1, t_2; \alpha, \beta)$ is a product of two univariate functions, see (3.3). We normalize the first factor $f_1(t_1)$ by the condition $f_1(\eta_{l_1}) = 1$. Due to the definition of η_{l_1} we have $F(\eta_{l_1}, 1 - \eta_{l_1}; \alpha, \beta) = (-1)^{l_2} e^{w_n(\eta_{l_1}; \alpha, \beta)} = 1$. Thus

$$F(t_1, t_2; \alpha, \beta) = f_1(t_1)f_2(t_2), \quad f_2(1 - \eta_{l_1}) = 1 (= f_1(\eta_{l_1})).$$

In addition, since $\xi_{l_1} < \eta_{l_1}$ we can easily check, see (1.4) and (3.4), that $f_1(t)$ is strictly increasing for $t \in [\eta_{l_1}, 1]$, in particular $f_1(t) \geq f_1(\eta_{l_1}) = 1$ here. Since $1 - \xi_{l_1+1} < 1 - \eta_{l_1}$, $f_2(t)$ is strictly increasing and $f_2(t) \geq 1$ for $t \in [1 - \eta_{l_1}, 1]$.

We note that by (3.4) and (1.3):

$$|F(t, 1 - t; \alpha, \beta)| = t^\alpha (1 - t)^\beta |J_{l_1+l_2}(t; \alpha, \beta)| \leq 1, \quad (3.5)$$

for all $t \in [0, 1]$. In order to show the main claim

$$|F(t_1, t_2; \alpha, \beta)| \leq 1 \quad (3.6)$$

for all $(t_1, t_2) \in \Delta$, we distinguish three regions in Δ .

If $t \in [0, \eta_{l_1}]$ then (3.5) yields $|f_1(t)| \leq 1/|f_2(1 - t)| \leq 1$, where the last inequality follows by the above listed properties of f_2 . Similarly we obtain $|f_2(t)| \leq 1$, if $t \in [0, 1 - \eta_{l_1}]$. Thus

$$|F(t_1, t_2; \alpha, \beta)| = |f_1(t_1)f_2(t_2)| \leq 1 \quad \text{for } t_1 \in [0, \eta_{l_1}], t_2 \in [0, 1 - \eta_{l_1}].$$

If $\eta_{l_1} \leq t_1 \leq 1 - t_2 \leq 1$, then since f_1 is increasing on $[\eta_{l_1}, 1]$, it follows that $|f_1(t_1)f_2(t_2)| \leq |f_1(1 - t_2)f_2(t_2)| = |F(1 - t_2, t_2; \alpha, \beta)|$, hence (3.6) follows by (3.5).

If $1 - \eta_{l_1} \leq t_2 \leq 1 - t_1 \leq 1$, then since f_2 is increasing on $[1 - \eta_{l_1}, 1]$, it follows that $|f_1(t_1)f_2(t_2)| \leq |f_1(t_1)f_2(1 - t_1)| = |F(t_1, 1 - t_1; \alpha, \beta)|$, hence (3.6) follows again by (3.5).

By combining the three cases it follows that relation (3.6) holds for all $(t_1, t_2) \in \Delta$.

In conclusion, relations (3.4) and (3.6) being proved, by Proposition 2.2 it follows that $P_{l_1, l_2}(t_1, t_2; \alpha, \beta)$ is a normalized polynomial of least deviation from zero on Δ with respect to the weight $t_1^\alpha t_2^\beta$, and hence, the polynomial given by (1.5) is a polynomial of least deviation from zero on $\mathbb{B}$, which also proves Theorem 1.1. $\square$




Figure 3: Domain $\Omega_*(\alpha, \beta)$

4 Leading term in asymptotics

We need certain properties of the conformal mapping w_* of the upper half-plane onto the domain

$$\Omega_* = \{w = u + iv : -\beta\pi < v < (1 - \beta)\pi\} \setminus \{w = u + iv : u \leq 0, 0 \leq v \leq \alpha\pi\},$$

see Fig. 3. Due to the Schwarz-Christoffel formula [5], it is of the form

$$w_*(z; \alpha, \beta) = \int_{x_2}^z \frac{\sqrt{(z - x_1)(z - x_2)}}{z(z - 1)} dz$$

where x_1, x_2 , $0 < x_1 < x_2 < 1$, are the preimages of the angle-points $\pi\alpha i$ and 0 respectively. As before three "infinite points" in the domain correspond to $0, 1$ and ∞ and due to the size of corresponding strips we have the following relations

$$\alpha = \sqrt{x_1 x_2}, \quad \beta = \sqrt{(1 - x_1)(1 - x_2)}. \quad (4.1)$$

This is an elementary integral, so we get

$$\begin{aligned} w_*(z; \alpha, \beta) = & \sqrt{x_1 x_2} \ln \frac{(\sqrt{x_2(1 - \frac{x_1}{z})} - \sqrt{x_1(1 - \frac{x_2}{z})})^2}{x_2 - x_1} \\ & + \sqrt{(1 - x_1)(1 - x_2)} \ln \frac{(\sqrt{(1 - x_2)(1 - \frac{1-x_1}{1-z})} - \sqrt{(1 - x_1)(1 - \frac{1-x_2}{1-z})})^2}{x_1 - x_2} \\ & + \ln \frac{(\sqrt{z - x_1} + \sqrt{z - x_2})^2}{x_2 - x_1}. \end{aligned} \quad (4.2)$$

Similarly to (1.1) we define the real constant $C_*(\alpha, \beta)$ by the condition

$$w_*(z; \alpha, \beta) = \ln z + C_*(\alpha, \beta) - \beta\pi i + O(1/z), \quad z \rightarrow \infty.$$

By (4.2) we get

$$\begin{aligned} C_*(\alpha, \beta) = & \sqrt{x_1 x_2} \ln \frac{(\sqrt{x_2} - \sqrt{x_1})^2}{x_2 - x_1} \\ & + \sqrt{(1-x_1)(1-x_2)} \ln \frac{(\sqrt{1-x_2} - \sqrt{1-x_1})^2}{x_2 - x_1} \\ & + \ln \frac{4}{x_2 - x_1}, \end{aligned}$$

which we simplify to

$$\begin{aligned} C_*(\alpha, \beta) = & (1 - \sqrt{x_1 x_2} - \sqrt{(1-x_1)(1-x_2)}) \ln \frac{4}{x_2 - x_1} \\ & + 2\sqrt{x_1 x_2} \ln \frac{2}{\sqrt{x_2} + \sqrt{x_1}} + 2\sqrt{(1-x_1)(1-x_2)} \ln \frac{2}{\sqrt{1-x_2} + \sqrt{1-x_1}}. \end{aligned}$$

Using (4.1) we get

$$\begin{aligned} C_*(\alpha, \beta) = & \frac{1-\alpha-\beta}{2} \ln \frac{16}{(1-(\alpha+\beta)^2)(1-(\alpha-\beta)^2)} \\ & + \alpha \ln \frac{4}{(1+\alpha)^2 - \beta^2} + \beta \ln \frac{4}{(1+\beta)^2 - \alpha^2}. \end{aligned} \quad (4.3)$$

Proof of Theorem 1.4. As the sequence of domains $\frac{2}{n}\Omega_{l_1+l_2}(\frac{k_1-l_1}{2}, \frac{k_2-l_2}{2})$ converges to Ω_* as $n \rightarrow \infty$, it follows by Carathéodory's theorem, see e.g. [7], that for the sequence of conformal mappings it holds that

$$w_*(z; \varkappa_1 - \lambda_1, \varkappa_2 - \lambda_2) = \lim_{n \rightarrow \infty} \frac{2}{n} w_{l_1+l_2}(z; \frac{k_1-l_1}{2}, \frac{k_2-l_2}{2})$$

Therefore

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{2}{n} \ln \Lambda_{k_1, l_1, k_2, l_2} &= \lim_{n \rightarrow \infty} \frac{2}{n} C_{l_1+l_2} \left(\frac{k_1-l_1}{2}, \frac{k_2-l_2}{2} \right) \\ &= C_*(\varkappa_1 - \lambda_1, \varkappa_2 - \lambda_2). \end{aligned}$$

Since in this case

$$\begin{aligned} 1 - \alpha - \beta &= 2(\lambda_1 + \lambda_2) \\ 1 + \alpha + \beta &= 2(\varkappa_1 + \varkappa_2) \\ 1 - \alpha + \beta &= 2(\lambda_1 + \varkappa_2) \\ 1 + \alpha - \beta &= 2(\varkappa_1 + \lambda_2) \end{aligned}$$

by (4.3) we get

$$\begin{aligned} C_*(\varkappa_1 - \lambda_1, \varkappa_2 - \lambda_2) = & -(\varkappa_1 + \varkappa_2) \ln(\varkappa_1 + \varkappa_2) \\ & -(\lambda_1 + \lambda_2) \ln(\lambda_1 + \lambda_2) \\ & -(\varkappa_1 + \lambda_2) \ln(\varkappa_1 + \lambda_2) \\ & -(\lambda_1 + \varkappa_2) \ln(\lambda_1 + \varkappa_2) \end{aligned}$$

□

References

- [1] N. I. Akhiezer, Лекции по теории аппроксимации. (Russian) [Lectures in the theory of approximation] Second, revised and enlarged edition Izdat. “Nauka”, Moscow 1965. 407 pp.
- [2] N. I. Akhiezer and B. Ya. Levin, *Generalization of S. N. Bernstein’s inequality for derivatives of entire functions*. (Russian) 1960 Issledovaniya po sovremennym problemam teorii funktsii kompleksnogo peremennogo pp. 111–165 Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow.
- [3] G. Andrews, R. Askey, R. Roy, *Special functions*. Encyclopedia of Mathematics and its Applications, 71. Cambridge University Press, Cambridge, 1999. 664 pp.
- [4] H. Braß, *Ein Beispiel zur Theorie der besten Approximation*. Multivariate Approximation Theory, II (Oberwolfach, 1982), Internat. Ser. Numer. Math. 61, Birkhäuser, Basel-Boston, Mass. 1982, 59-67.
- [5] T. A. Driscoll and L. N. Trefethen, *Schwarz-Christoffel mapping*. Cambridge Monographs on Applied and Computational Mathematics, 8. Cambridge University Press, Cambridge, 2002. 132 pp.
- [6] A. Eremenko and P. Yuditskii, *Uniform approximation of $\operatorname{sgn} x$ by polynomials and entire functions*. J. Anal. Math. 101 (2007), 313–324.
- [7] G. M. Galuzin, *Geometric Theory of Functions of a Complex Variable*, Amer. Math. Soc. Providence.(1969), Vol. 26.
- [8] W. B. Gearhart, *Some Chebyshev approximations by polynomials in two variables*. J. Approx. Theory 8 (1973), 195-209.

- [9] B. Ya. Levin, *Classification of closed sets on R and representation of a majorant*. III. (Russian) Teor. Funktsii Funktsional. Anal. i Prilozhen. No. 52 (1989), 21–33; translation in J. Soviet Math. 52 (1990), no. 5, 3364–3372.
- [10] G. MacLane, *Concerning the uniformization of certain Riemann surfaces allied to the inverse-cosine and inverse-gamma surfaces*, Trans. Amer. Math. Soc. 62 (1947), 99–113.
- [11] I. Moale and F. Peherstorfer, *Explicit min-max polynomials on the disc*. Submitted.
- [12] F. Nazarov, F. Peherstorfer, A. Volberg, P. Yuditskii, *Asymptotics of the best polynomial approximation of $|x|^p$ and of the best Laurent polynomial approximation of $\operatorname{sgn}(x)$ on two symmetric intervals*. Constr. Approx. 29 (2009), no. 1, 23–39.
- [13] M. Reimer, *On multivariate polynomials of least deviation from zero on the unit ball*. Math. Z. 153 (1977), 51–58.
- [14] M. Sodin and P. Yuditskii, *Functions that deviate least from zero on closed subsets of the real axis*. (Russian) Algebra i Analiz 4 (1992), no. 2, 1–61; translation in St. Petersburg Math. J. 4 (1993), no. 2, 201–249.
- [15] E. B. Vinberg, *Real entire functions with prescribed critical values*, in Problems of Group Theory and Homological Algebra, Yaroslavl. Gos. U., Yaroslavl, 1989, pp. 127–138.

*Institute for Analysis, Johannes Kepler University,
A-4040 Linz, Austria*

E-mail: Ionela.Moale@jku.at

E-mail: Petro.Yudytskiy@jku.at